

**Brandon M. Tauberg, M.D.**  
 Orthopaedic Sports Medicine Surgeon  
 Arthroscopic and Shoulder Surgery  
 www.brandontaubergmd.com

**South Hills Location**  
 2000 Oxford Drive  
 Suite 211, 2<sup>nd</sup> Floor  
 Pittsburgh, PA 15102  
 412-283-0260 (office)  
 412-283-0070 (fax)

## *Meniscal Repair Protocol*

	WEIGHT BEARING	BRACE	ROM	EXERCISES	PROGRESSION GOALS
<b>PHASE I - Acute</b> <b>0-6 weeks</b>	<ul style="list-style-type: none"> <li>• 0-2 Weeks:               <ul style="list-style-type: none"> <li>- TTWB</li> </ul> </li> <li>• 2-4 Weeks:               <ul style="list-style-type: none"> <li>- PWB</li> </ul> </li> <li>• 4+ Weeks:               <ul style="list-style-type: none"> <li>- FWB with symmetrical gait</li> </ul> </li> <li>• Unlock brace when adequate quad control</li> <li>• Wean off crutches as gait normalizes</li> </ul>	<ul style="list-style-type: none"> <li>• 0-1 Week:               <ul style="list-style-type: none"> <li>- Brace locked</li> <li>0° extension</li> </ul> </li> <li>• 1-2 Weeks:               <ul style="list-style-type: none"> <li>- Unlock brace</li> <li>0° to 60°</li> </ul> </li> <li>• 2-4 Weeks:               <ul style="list-style-type: none"> <li>- Unlock brace</li> <li>0° to 90°</li> </ul> </li> <li>• 4-6 Weeks:               <ul style="list-style-type: none"> <li>- Brace unlocked to full if good quad control and full extension</li> </ul> </li> <li>• 6+ Weeks:               <ul style="list-style-type: none"> <li>- D/C crutches when gait is normal</li> <li>- D/C brace for home ambulation</li> <li>- Continue brace for community</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• 0-1 Week:               <ul style="list-style-type: none"> <li>- Full extension and progress flexion to 60°</li> </ul> </li> <li>• 1-4 Weeks:               <ul style="list-style-type: none"> <li>- Maintain full extension and progress to 90°</li> </ul> </li> <li>• 4+ Weeks:               <ul style="list-style-type: none"> <li>- Gradually progress flexion to 120°</li> </ul> </li> </ul> <p><b>STRESS EARLY EXTENSION</b> (avoid hyperextension &gt; 10°)</p>	<ul style="list-style-type: none"> <li>• Quad sets, straight leg raises, hip abduction, patellar mobs, SAQ*</li> <li>• 2+ weeks               <ul style="list-style-type: none"> <li>- TKEs, calf raises, pre-gait</li> </ul> </li> <li>• 4+ weeks               <ul style="list-style-type: none"> <li>- Static balance</li> <li>- Mini squats</li> <li>- Stationary biking: must be &gt;110° knee flexion</li> </ul> </li> <li>• No weight bearing with flexion &gt;90°</li> </ul>	<p><b>Criteria for Full Ambulation:</b></p> <ul style="list-style-type: none"> <li>• Knee extension ≥ 0°</li> <li>• Knee Flexion 90°</li> <li>• Minimal effusion/pain</li> <li>• Symmetrical gait without limp</li> <li>• MD or PT APPROVAL</li> </ul>

\* Avoid any tibial rotation for 8 weeks to protect meniscus



**South Hills Orthopaedic**

SURGERY ASSOCIATES, P.C.

## Meniscal Repair Protocol

	WEIGHT BEARING	BRACE	ROM	EXERCISES	PROGRESSION GOALS
		ambulation until MD approval to D/C			
<b>PHASE II - Strength</b> <b>6-12 weeks</b>	<ul style="list-style-type: none"> <li>As tolerated</li> </ul>	<ul style="list-style-type: none"> <li>Discontinued at 6 weeks if no extension lag</li> <li>Unlock and wean out of/off crutches at 6-8 weeks</li> </ul>	<ul style="list-style-type: none"> <li>6+ Weeks:                             <ul style="list-style-type: none"> <li>Maintain full extension</li> <li>Progress to full flexion</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>Continue with Phase I</li> <li>Strengthening                             <ul style="list-style-type: none"> <li>Loaded flexion &gt;90° is <b>PROHIBITED</b></li> <li>Leg press, step ups, step downs, RDLs, lunges, Bulgarian squats, wall sits</li> <li>Squat progression: bodyweight → single leg</li> <li>Advance hip abduction &amp; glute strength: band walks, lateral lunge, reverse lunge, bridges, hip thrusters</li> <li>Core exercises: planks, side planks, v-ups, Russian twist, superman</li> <li>Balance training: foam pad, balance board, BOSU</li> </ul> </li> <li>Conditioning                             <ul style="list-style-type: none"> <li>Initiate dynamic warm-up: Frankenstein kicks, leg swings, knee hugs, heel sweeps, heel/toe walks, oil rigs, lateral lunge, hip rotation, inch worm, speed squats</li> <li>Stationary bike</li> </ul> </li> </ul>	<b>Criteria for Phase 2:</b> <ul style="list-style-type: none"> <li>Pain &lt; 3/10 (worst)</li> <li>Within 2° normal knee extension &amp; 120° knee flexion</li> <li>Symmetrical body weight squat</li> <li>Minimal effusion</li> <li>Minimal pain</li> <li>Symmetrical gait without a limp</li> </ul>
<b>PHASE III – Initiate Jogging and Double Leg Plyometrics</b> <b>12-20 weeks</b>	<ul style="list-style-type: none"> <li>Full</li> </ul>	<ul style="list-style-type: none"> <li>None</li> </ul>	<ul style="list-style-type: none"> <li>12+ Weeks:                             <ul style="list-style-type: none"> <li>Maintain symmetry &amp; pain-free with overpressure</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>Strengthening                             <ul style="list-style-type: none"> <li>Leg press, step ups, step downs, RDLs, lunges, Bulgarian squats, wall sits</li> <li>Squat progression: bodyweight → single leg</li> <li>Advance hip abduction &amp; glut strength: band walks, lateral lunge, reverse lunge, bridges, hip thrusters</li> <li>Core exercises: planks, side planks, v-ups, Russian twist, superman</li> <li>Balance training: foam pad, balance board, BOSU</li> </ul> </li> <li>Conditioning                             <ul style="list-style-type: none"> <li>Dynamic warmup &amp; integrate sport specific warmup</li> <li>Stationary bike, elliptical, swimming (14 weeks), &amp; rowing machine</li> <li>12+ Weeks: treadmill walk/jog progressions. Begin with 30"-1' W/J intervals, advance jog time by 1 min each week</li> <li>12+ Weeks: Double leg jump rope and shuttle jumps</li> <li>14+ Weeks: Swimming → progress kicking gradually and pain-free, no flip turns</li> </ul> </li> </ul>	<b>Criteria For Jogging &amp; Double Leg Jump Rope</b> <ul style="list-style-type: none"> <li>Pain ≤ 3/10 (worst)</li> <li>Within 2° normal knee extension &amp; 120° knee flexion</li> <li>Quad and hamstring strength ≥ 60% normal</li> <li>Less than 4cm deficit on single-leg squat (anterior reach)</li> <li>≥ 1 minute single leg squats</li> <li>MD approval</li> </ul>

\* Avoid any tibial rotation for 8 weeks to protect meniscus



# Meniscal Repair Protocol

	WEIGHT BEARING	BRACE	ROM	EXERCISES	PROGRESSION GOALS
				<ul style="list-style-type: none"> <li>- 16+ Weeks: Advance to track workouts: jog straights and walk curves</li> </ul>	
<b>PHASE IV – Strength, Agility, Plyometrics</b> <b>20-24 weeks</b>	<ul style="list-style-type: none"> <li>• Full</li> </ul>	<ul style="list-style-type: none"> <li>• Functional bracing dependent on patient activity and doctor recommendation</li> </ul>	<ul style="list-style-type: none"> <li>• 20+ weeks: promote and maintain symmetry</li> </ul>	<ul style="list-style-type: none"> <li>• <b>Strengthening</b> <ul style="list-style-type: none"> <li>- Gym strengthening: squats, deadlifts, initiate Olympic lifting</li> <li>- Dynamic eccentric loading: double &amp; single leg</li> <li>- Dynamic core: rotational and anti-rotational drills</li> <li>- Integrate interval strength circuits &amp; work/rest timed intervals</li> <li>- Isokinetic training protocols: begin with 300°/sec, progress to 180°/sec</li> </ul> </li> <li>• <b>Conditioning</b> <ul style="list-style-type: none"> <li>- Dynamic warmup &amp; sport specific warmup</li> <li>- Stationary bike, elliptical, swimming, &amp; rowing machine</li> <li>- Track workouts: advance to linear speed drills and sprinting drills</li> </ul> </li> <li>• <b>Plyometrics &amp; Agility</b> <ul style="list-style-type: none"> <li>- Ladder drills, footwork agility drills, cone drills</li> <li>- Double leg plyos: jump rope, line jumps, cone jumps, depth jumps, box jumps</li> <li>- Single leg landings: alternating, line jumps, hops → SL jumps</li> <li>- High intensity predictable patterned movements, incorporate sport specific drills</li> <li>- Change of direction drills: begin with &lt;90°, progress to 90° and greater</li> </ul> </li> <li>• <b>Advanced Agility &amp; Plyometrics (2-3 days/week)</b> <ul style="list-style-type: none"> <li>- Tuck jumps, squat jumps, bounding, SL hop, SL triple hop, SL cross over hop</li> <li>- Change of direction drills</li> <li>- Introduce unpredictable agility movements</li> <li>- Non-contact sport-specific drills</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>• <b>Criteria for Plyometrics &amp; Agility:</b> <ul style="list-style-type: none"> <li>• Pain ≤ 2/10 (Worst)</li> <li>• Quad &amp; HS symmetry ≥ 80% normal; ≥ 50% H/Q ratio for females</li> <li>• ≤5 on landing error scoring system (LESS)</li> <li>• At least 3 minutes of single-leg squats (resisted)</li> <li>• Jogging &gt;15 minutes on treadmill</li> <li>• MD or PT APPROVAL</li> </ul> </li> <li>• <b>Criteria for Advanced Agility &amp; SL Plyometrics:</b> <ul style="list-style-type: none"> <li>• Pain &lt; 2/10 (Worst)</li> <li>• Quad &amp; HS symmetry ≥ 80% normal; ≥ 50% H/Q ratio for females</li> <li>• At least 3 minutes of single-leg squats (resisted)</li> <li>• Jogging &gt;15 minutes on track or paved surface</li> <li>• MD or PT APPROVAL</li> </ul> </li> </ul>



## Meniscal Repair Protocol

	WEIGHT BEARING	BRACE	ROM	EXERCISES	PROGRESSION GOALS
<b>PHASE VI – Return to Play 24+ weeks</b>	<ul style="list-style-type: none"> <li>• Full</li> </ul>	<ul style="list-style-type: none"> <li>• Functional bracing dependent on patient activity and doctor recs</li> </ul>	<ul style="list-style-type: none"> <li>• Full</li> </ul>	<ul style="list-style-type: none"> <li>• Strengthening                             <ul style="list-style-type: none"> <li>- Gym strengthening: squats, deadlifts, initiate Olympic lifting</li> <li>- Interval strength circuits &amp; work/rest timed intervals</li> <li>- Dynamic strength and core exercises</li> <li>- Complex movement patterns</li> <li>- Isokinetic protocols: 300°, 180°, 60°/sec</li> </ul> </li> <li>• Conditioning                             <ul style="list-style-type: none"> <li>- Biking, jogging, swimming, rowing, interval sprints</li> </ul> </li> <li>• Plyometrics &amp; Agility (2-3 days/week)                             <ul style="list-style-type: none"> <li>- Max effort DL and SL jumps → progress with rotation</li> <li>- Lateral &amp; rotational agility drills</li> <li>- Unpredictable cutting agility</li> <li>- Non-contact drills → contact drills with MD approval</li> <li>- Return to practice → contact practice → scrimmage → interval play → full play</li> </ul> </li> </ul>	<b>Return to Play Criteria:</b> <ul style="list-style-type: none"> <li>• VAS &lt; 2 ( Worst)</li> <li>• &gt;75/100 on ACL-RSI survey</li> <li>• Quad &amp; Hamstring strength ≥ 90% normal; ≥ 60% H/Q ratio for females</li> <li>• 90% normal on single-leg hop tests</li> <li>• 95% normal figure of 8, SL vertical jump, &amp; 5-10-5 pro-agility</li> <li>• MD APPROVAL</li> </ul>

- Do NOT change bandages unless instructed by physician
- Monitor for pain and swelling. Modify as necessary.
- Encourage home exercises program daily
- Encourage ice 4x a day for 20 minutes while swelling is present.
- For any questions or concerns please contact Dr. Tauberg's office



1. Surgical pre-cautions: Do not change bandages unless instructed by physician. If you suspect a DVT, contact Dr. Tauberg's office immediately at 412-283-0260 or refer to ED immediately. If patient has reactive effusion that does not improve with rest, ice, and compression, contact Dr. Tauberg's office.
2. Begin stretching extension ROM on day one. Achieve full extension ROM by week 2. If not achieved by end of week 4, notify the physician's office.
3. Address quad activation early and focus on isolation of quadriceps activation. Use surface EMG, NMES, and tactile cueing to isolate quadriceps. Be aware of co-contracting from hamstrings, and ensure proper form. Do not progress to standing activities if patient is unable to achieve isolated quad set in long seated position. Goal is to achieve heel lift with a quad set. \*Dosing quad sets: 10 minutes of 10 second squeeze/10 second rest, x5 times a day.
4. Straight leg raises: Ensure quadriceps is activated and is maintaining contraction throughout the SLR range to eliminate extensor lag. Aim for a calf tap and elimination of extensor lag by week 3. Calf tap: the calf taps/skims the table while the heel stays elevated as the leg descends to starting position. Continue doing SLR until 10# ankle weight is achieved.
5. Do not force **flexion ROM**, but encourage steady progression. Patellar mobility is imperative. Use gentle soft tissue techniques for areas such as anterior interval/fat pad, quadriceps, hamstrings, and scar management. If 90° of flexion is not achieved by week 4, notify physician's office.
6. Start double leg (DL) mini squats and leg press from 0° to 60° initially, then progress to 90° as tolerated. Single leg (SL) activities may be initiated at week 4 with SL leg press and step-ups, then advancing to SL activities as tolerated. Loaded leg extensions are prohibited.  
\*Squat progressions example: DL leg press, DL mini squats, DL chair squats, DL body weight squats, SL leg press, SL step ups, Static lunge split squat, SL step downs, SL squats, SL split squat with elevated back leg, walking lunges, SL sit to stands, SL slide outs.
7. Pre-run/pre-jump program includes tempo-based activities with focus on the deceleration phase such as DL speed squats, DL drop squats, DL "bounce bounce squat", then progress to alternating SL drop squats. Also, intermittently increase the tempo of regular strengthening exercises to align with the timing requirements of jogging and jumping.
8. Walk/Jog program: MD approval required. Begin on treadmill with 2- 3 days per week. Begin with 1:1 or 2:1 walk to jog ratios, (i.e. 1 min walk to 1 min jog or 2 min walk to 1 min jog). Then progress each week by 1 min jog until 12-15 min of jogging is achieved.
9. Plyometric program: MD approval required. Begin with small DL jumps, jump rope, and small depth jump landings& box jumps. Progress box height as skill is mastered. Ensure equal weighted DL take-off and landing before progressing to SL plyometrics. Initiate SL plyometrics with alternating L and R landings in place and then advance to SL hops. Begin a sports metric based plyometric program when released by surgeon.
10. Isokinetic protocol: After 16 weeks and with MD approval, may begin training and testing with 300°/sec and progress to 180°/sec. Do not proceed if patient has history of anterior knee pain.
11. Return to Play Progression: a graded re-exposure is essential. Return to non-contact practice, return to contact practice, return to scrimmage, return to interval play, return to full time play

